



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

some specimens, measuring 2.94x2.16 and 2.90x2.16 inches. Both of them show traces of faint vinaceous and vinaceous-cinnamon markings scattered at random over the surface. In this respect they are of the very rarest type, but I might here add that one other set, taken by the same gentleman, shows these faint markings. These eggs are also very large specimens, measuring 3.08x2.18 and 2.92x2.18 inches.

J. WARREN JACOBS, *Waynesburg, Pa.*

BIRD NOTES FROM SOUTH-EASTERN IOWA: BLUE GOOSE (*Chen caerulescens*).—While hunting, March 21, 1908, about Green Bay, Lee County, Iowa, we found the Blue Goose. I think that day and the day before they were fairly common there; and we saw six together and shot one. I think they are quite rare, however, for old hunters could hardly believe there were any about.

RED-HEADED WOODPECKER (*Melanerpes erythrocephalus*).—This bird was very common, in fact, abundant, in a large woods near Green Bay, March 19-21, when we were there. This is only six miles from here (Denmark, Iowa), yet none did I see here until April 22.

LECONTE'S SPARROW (*Ammodramus lecontei*).—This elusive sparrow was very common in the dried weeds near Green Bay, March 19-21. We frequently found them in flocks of half a dozen or more. They were exceedingly difficult to see and I shot two finally to make sure of the identification. A good many times I watched them drop down into a bunch of weeds and would literally have to kick the bunch before they would fly out, only to drop into the next one. They were fairly common here about Denmark (if one knew where to look for them) till April 21. Nearly every time I have found Swamp Sparrows in company with them.

ROBERT L. BAIRD, *Denmark, Iowa.*

BRUNNICH'S MURRE ON LAKE ERIE, 1907.—A Brunnich's Murre was found in the eastern part of Sandusky, Ohio, in the water near Big Island, and picked up December 1st, 1907. It lived two or three days in captivity. I had it mounted. The taxidermist told me lately that two others were brought to him at the same time. One of my pupils told me that four of these birds were shot December 3d, by the Teasel Brothers, who said they could have shot two more. It may be that the two not killed were those mentioned by the taxidermist.

E. L. MOSELEY, *Sandusky, Ohio.*